ADHD Treatment Can Make a Real Difference



WHAT IS DYANAVEL XR?

DYANAVEL XR (amphetamine) is a central nervous system (CNS) stimulant prescription medicine used for the treatment of Attention Deficit Hyperactivity Disorder (ADHD) in people 6 years and older. **DYANAVEL XR may help increase attention and decrease impulsiveness and hyperactivity in people with ADHD.**

It is not known if DYANAVEL XR is safe and effective in children under 6 years of age.

IMPORTANT SAFETY INFORMATION

DYANAVEL XR is a federally controlled substance (CII) because it can be abused or lead to dependence. Keep DYANAVEL XR in a safe place to prevent misuse and abuse. Selling or giving away DYANAVEL XR may harm others, and is against the law. Tell your physician if you, your child, or any family members have ever abused or been dependent on alcohol, prescription medicines, or street drugs.

Please see additional Important Safety Information throughout, and Medication Guide and Full Prescribing Information, including Boxed Warning regarding Abuse and Dependence.

Next Page

What Is ADHD?

Attention Deficit Hyperactivity Disorder, or ADHD, is a disorder that affects the structure and function of the brain

The exact causes and risk factors for ADHD are unknown, but it is thought that genetics may play an important role. In 41% to 55% of families with a child who has ADHD, at least one parent is also thought to be affected.

ADHD in the **United States**

ages 18 to 44

ages 2 to 17

Symptoms of ADHD

People with ADHD experience a pattern or combination of three types of symptoms that can interfere with different aspects of life.



> Inattention

A person may have a hard time staying on task, remaining focused, or keeping organized.



Hyperactivity

A person may seem to move about constantly, including when it is not appropriate. This could also mean a person excessively fidgets, taps, or talks too much.



Impulsivity

A person may act without thinking or have difficulty with self-control.

ADHD may look different in different people. Only a healthcare professional can diagnose ADHD, so it's important to talk to your doctor about your symptoms

ADHD Can Be Disruptive

ADHD symptoms can interfere with aspects of your life such as work, school, relationships, or social interaction

ADHD can affect many day-to-day activities depending on your unique situation, including some of the following:





Morning

- Organizing yourself or your kids for the day
- Getting to work or school on time





Afternoon

- Paying attention at work or school
- Finishing work accurately and on time
- Helping kids with homework





- Interacting with family or friends
- Managing finances
- Planning the next day

About 60% of people with ADHD have symptoms that persist from childhood into adulthood

Find an ADHD Treatment Option That Works for You

Your healthcare provider will work with you to determine a treatment plan that helps you manage your ADHD symptoms

While there is currently no cure for ADHD, there are several options available to reduce symptoms and help improve functioning. These include medications, psychotherapy, education or training, or a combination of treatments.

Medication for ADHD typically comes in the form of stimulants or nonstimulants

Stimulants are the most widely used ADHD medications

Nonstimulants may be used when stimulants do not work well, have unacceptable side effects, or a nonstimulant is preferred for other reasons

Nonmedication options are often used alongside medication

- Cognitive behavioral therapy helps you develop a more mindful and structured approach to thinking and behaving
- Psychoeducation can include a coach who helps you develop life skills such as planning, management, and self-esteem
- College and work accommodations can be developed for individual situations

For more information about these options, talk to your healthcare provider.



It Is Important to Treat ADHD

When deciding whether or not to seek treatment, consider the areas of your life that could be impacted



Work or school life

Difficulty with attention, time management, and staying on task



Social interactions

Lack of impulse control that can result in angry outbursts

Daily activities

Problems with self-organization that can make it difficult to follow a routine



With the right treatment, ADHD can be managed and symptoms may improve

DYANAVEL XR Tablet May Help Manage Your **ADHD Symptoms**

For patients 6 years and older with ADHD, DYANAVEL® XR (amphetamine) tablet may help increase attention and decrease impulsiveness and hyperactivity

DYANAVEL XR is a central nervous system stimulant prescription medicine used for the treatment of ADHD.

It is not known if DYANAVEL XR is safe and effective in children under 6 years of age.



Innovation

DYANAVEL XR tablet works through a technology called LiquiXR® (pronounced li-quee-x-r). Despite its name, this technology is used to develop tablet medications by combining fast-acting and long-lasting components all in one.

LiquiXR uses a specific process to release medication continuously throughout the day without any sudden drops in medication levels.



(&) Safety and effectiveness

Studied in adults with ADHD aged 18 to 60 years.



Dosing

DYANAVEL XR tablet has the most dosing options of any extended-release amphetamine tablet, so your doctor can work with you to find a personalized dose that helps control symptoms and reduces side effects. DYANAVEL XR tablet should be taken exactly as prescribed by your doctor. Talk to your doctor about side effects that bother you or do not go away.

IMPORTANT SAFETY INFORMATION (cont'd)

DYANAVEL XR should not be taken if you or your child are allergic to amphetamine or any of the ingredients in DYANAVEL XR, or are taking or have taken within the past 14 days an anti-depression medicine called a monoamine oxidase inhibitor or MAOL

DYANAVEL XR can cause serious side effects. Tell the doctor:

 if you or your child have heart problems, heart defects, high blood pressure, or a family history of these problems. This is important because sudden death has occurred in children 6 to 17 years old with heart problems or defects, and sudden death, stroke, and heart attack have happened in adults. Your doctor should check you or your child carefully for heart problems before starting DYANAVEL XR. Since increases in blood pressure and heart rate may occur, the doctor should regularly check these during treatment. Call the doctor right away if you or your child have any signs of heart problems such as chest pain, shortness of breath, or fainting while taking DYANAVEL XR.



DYANAVEL XR (amphetamine) tablet

Your Doctor Can Find the Right Dose for You



IMPORTANT SAFETY INFORMATION (cont'd)

DYANAVEL XR (amphetamine) can cause serious side effects. Tell the doctor:

- if you or your child have mental problems, or a family history of suicide, bipolar illness, or depression. This is important because new or worsening behavior and thought problems or bipolar illness may occur. New symptoms such as seeing or hearing things that are not real, believing things that are not true, being suspicious, or having new manic symptoms may occur. **Call the doctor right away if there are any new or worsening mental symptoms during treatment.**
- if you or your child have circulation problems in fingers and toes (called peripheral vasculopathy, including Raynaud's phenomenon). Fingers or toes may feel numb, cool, painful, sensitive to temperature, and/ or change color from pale, to blue, to red. **Call the doctor right away if any signs of unexplained wounds appear on fingers or toes while taking DYANAVEL XR.**





Tris Pharma Offers Support to Help You Get Started

Co-pay savings are available for eligible patients. Terms and Conditions apply.



Ask your doctor for a co-pay card.

You can also get a mobile savings card,* along with helpful refill reminders, by texting TRIS to TRIS4U (874748).

Access resources to learn more about ADHD and support available to you

Visit coupon.trisadhd.com to find:

- Community resources for ADHD
- Helpful information about Tris ADHD products
- A pharmacy locator and much more

IMPORTANT SAFETY INFORMATION (cont'd)

DYANAVEL XR (amphetamine) can cause serious side effects. Tell the doctor:

- if your child is having slowing of growth (height and weight); Your child should have his or her height and weight checked often while taking DYANAVEL XR.
- if you or your child have symptoms of serotonin syndrome agitation, hallucinations, coma, or other changes in mental status; fast heartbeat; sweating or fever; nausea, vomiting, diarrhea; high or low blood pressure; problems controlling movements or muscle twitching; muscle stiffness or tightness. Serotonin syndrome may occur when DYANAVEL XR is taken with certain other medicines and may be life-threatening. Stop DYANAVEL XR and call your doctor or go to the nearest hospital emergency room.

^{*}Mobile Terms and Conditions: coupon.trisadhd.com/terms. Message data rates may apply.

Additional Help and Support

Centers for Disease Control and Prevention (CDC) Attention-Deficit / Hyperactivity Disorder (ADHD)

cdc.gov/ncbddd/adhd

National Institute of Mental Health (NIMH)

nimh.nih.gov/health/topics/attention-deficit-hyperactivity-disorder-adhd

Children and Adults with Attention-Deficit/Hyperactivity Disorder (CHADD) chadd.org

HealthyChildren.org from the American Academy of Pediatrics healthychildren.org

Attention Deficit Disorder Association (ADDA) add.org

Tris Pharma Has a Family of Proven ADHD Products for Patients Aged 6 Years and Above

Why the need for options? No two people are alike. Why should their medicine, or their dose, be the same? Tris products cover both main types of stimulants for ADHD, methylphenidate and amphetamine. And the good news is that about 85% of patients with ADHD respond to one stimulant or the other.





IMPORTANT SAFETY INFORMATION (cont'd)

DYANAVEL XR (amphetamine) can cause serious side effects. Tell the doctor:

• if you or your child are pregnant or plan to become pregnant. It is not known if DYANAVEL XR will harm your unborn baby. Tell your doctor if you or your child become pregnant during treatment with DYANAVEL XR.

For more information about Tris ADHD products, please visit **trisadhd.com**





IMPORTANT SAFETY INFORMATION (cont'd)

DYANAVEL XR (amphetamine) can cause serious side effects. Tell the doctor:

• if you or your child is breastfeeding or plan to breastfeed. DYANAVEL XR passes into breast milk. You or your child should not breastfeed while you are taking DYANAVEL XR.

dizziness

• increased heart rate

Common side effects of amphetamine products include:

- dry mouth
- decreased appetite
- weight loss
- stomach pain
- nausea
- trouble sleeping
- restlessness
- restiessiless
- extreme mood changes

Talk to your doctor if you or your child have any side effects that bother you or do not go away.

Avoid drinking alcohol while taking DYANAVEL XR.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see Medication Guide and <u>Full Prescribing Information</u>, including Boxed Warning regarding Abuse and Dependence.

